

SUGAR-CONTAINING AMPHIPHILIC OLIGOMERS

FIG.1A

FIG.1B

FIG.1C





11







FIG.4







(SEQ ID NO: 1) ANALGESIC EFFECT OF A 5 mg/kg IV DOSE OF CETYL-PEG2 -ENKEPHALIN MONOCONJUGATE IN THE RAT HOT-PLATE ASSAY




FIG.7

COMPARISON OF μ -RECE	RECEPTOR BINDING AFFINITY OF ENKEPHALIN CONJUGATES	JUGATES
DRUG OR CONJUGATE	DETAILED STRUCTURE	% SPECIFIC BINDING
NALOXONE	NALOXONE	100
ENKEPHALIN	MET-ENKEPHALIN-LYS (SEQ ID NO: 49)	67
CETYL-ENK	CETYL-PEG ₂ -ENK (SEQ ID NO: 1)	100
CHOL-ENK	CHOLESTEROL-PEG3-ENK (SEQ ID NO: 1)	95
DHA-ENK	DHA-PEG ₂ -ENK (SEQ ID NO: 1)	63
PALM-ENK	PALMITATE-PEG3-ENK (SEQ ID NO: 1)	76
CETYL-TEG-ENK	CETYL-PEG3-ENK (SEQ ID NO: 1)	100

FIG.8

SYNTHESIS OF OLICOMER

Ŧ

BASE

\ \ \ \ \ m HYLENE GLYCOL =0, 1, 2, 3

ADDITION

Me-(CH₂)_nBr OMO ALKYL DERIVATIVE)

AMPHIPHILIC POLYMER

FREE END'

H_0 (0)

PIC 0

AMPHIPHILIC POLYMER

ATTACHMENT OF OLIGOMER TO ENKEPHALIN

OLIGOMER-ENKEPHALIN-CONJUGATE

m=14 AND n=2

CETYL-PEG2-ENKEPHALIN